Anatomical basis of diverse jaw protrusion directionality in ponyfishes (Family Leiognathidae)



Running title: Jaw protrusion diversity in ponyfishes

Alexus S. Roberts¹*, Jennifer R. Hodge², Prosanta Chakrabarty³, Peter C. Wainwright¹

 ¹Department of Evolution and Ecology, University of California Davis, Davis, CA 95616
²Department of Biological Sciences, Clemson University, Clemson, SC 29634
³Department of Biological Sciences, Museum of Natural Science, Louisiana State University, Baton Rouge, LA 70803

Contact Information: Alexus Roberts <u>asroberts@ucdavis.edu</u> Department of Evolution and Ecology University of California Davis 1 Shields Avenue Davis, CA 95616

Abstract: Protrusion of the oral jaws is a key morphological innovation that enhances feeding performance in fishes. The mechanisms of protrusion and the basis of variation in its magnitude are well studied, but little attention has been paid to the functional morphology of protrusion directionality, despite wide variation among teleost species from slightly dorsal to strongly ventral. Ponyfishes (Leiognathidae) comprise a group of 52 species that exhibit striking diversity in the directionality of jaw protrusion, providing a promising system for exploring its underlying basis in a single clade. We examined the anatomical basis of protrusion directionality by measuring eight traits associated with the size and positioning of oral jaw bones. Measurements were made on cleared and stained specimens of 20 ponyfish species, representing every major lineage within the family. Species fell into three non-overlapping clusters with respect to directionality including dorsal, rostral, and ventral protruders. A key correlate of protrusion direction is the anterior-posterior position of the articular-quadrate jaw joint. As the joint position moves from a posterior to a more anterior location, the orientation of the relaxed mandible

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/jmor.21314

rotates from an almost horizontal resting position to an upright vertical posture. Abduction of the mandible from the horizontal position results in ventrally directed protrusion, while the more upright mandible rotates to a position that maintains dorsal orientation. The resting orientation of the premaxilla and maxilla, thus, vary consistently with protrusion direction. Mouth size, represented by length of the mandible and maxilla, is a second major axis of variation in ponyfishes that is independent of variation in protrusion directionality.

Keywords: Slipmouths, Acanthomorpha, premaxillary protrusion, articular-quadrate jaw joint

Highlights:

- Ponyfishes (Leiognathidae) comprise a group of 52 acanthomorph species that exhibit striking diversity in the orientation of oral jaw protrusion (ranging more than 60°), with species that protrude their jaws in either a ventral, rostral, or dorsal direction.
- The size and orientation of craniofacial elements including the premaxillary ascending process, mandible, maxilla, and jaw joint position distinguish the three oral jaw functional groups.
- The position of the articular-quadrate jaw joint is a key correlate of protrusion directionality, as its movement along an anterior-posterior axis affects the resting and abducted orientation of the jaw bones.

Graphical Abstract: Leiognathids comprise a group of 52 acanthomorph species that exhibit striking craniofacial diversity. Direction of oral jaw protrusion, which correlates strongly with the anterior-posterior position of the articular-quadrate joint, represents a major axis of variation in this clade, with species that protrude their jaws in either a ventral (top left), rostral (bottom), or dorsal (top right) direction. Mouth size also represents a major axis of variation in ponyfishes that is independent of variation in protrusion directionality.



1. INTRODUCTION

The capacity for premaxillary protrusion is a major innovation in the jaws of fishes that results in several enhancements of feeding performance (Muller & Osse, 1978; Gosline, 1980; Motta, 1984). Jaw protrusion facilitates greater dexterity during various biting and manipulation behaviors (Konow & Bellwood, 2005; Gibb et al., 2015) and results in a substantial increase in suction feeding performance (Lauder, 1980; Liem, 1980; Holzman et al., 2007). During suction feeding, jaw protrusion plays a central role in rapidly positioning the mouth aperture close to the prey item, allowing the spatially restricted suction flow maximum exposure to the prey (Holzman et al., 2007). The capacity for premaxillary jaw protrusion has evolved several times in ray-finned fishes, including independent origins in two of the most successful radiations, Cypriniformes and Acanthomorpha (Wainwright et al., 2015).

While suction feeding movements in most fishes extend the mouth aperture in a rostral direction, numerous species that feed on benthic prey extend the mouth ventrally (Chao & Musick, 1977; Liem, 1979; Gosline, 1984), while fewer protrude somewhat dorsally and approach their prey from below (Alexander, 1967a; Liem, 1967; Pietsch, 1978; Grobecker, 1983). The diversity in protrusion directionality has been recognized for some time in the ecomorphological literature because it relates strongly to where fish feed, but the functional morphology that underlies this variation is poorly known.

Ponyfishes (Family Leiognathidae) comprise a group of over 50 bioluminescent acanthomorph species that show striking diversity in jaw protrusion directionality (Figure 1). Their common names, 'ponyfish' and 'slipmouth', refer to the horse-shaped cranial profile when the jaws are

protracted and the great extent of this protrusion, respectively (Nelson et al., 2016). These fishes are found in sandy-shore habitats and open waters of the Indo-West Pacific. They feed on small benthic organisms and midwater zooplankton, with diet varying among species (Jones, 1985; Jing, 1997). Although extensively studied to understand their circumesophageal light organ system (Harvey, 1921; Hastings, 1971; McFall-Ngai & Dunlap, 1984; Woodland et al., 2002; Sasaki et al., 2003; Chakrabarty et al., 2011a), less is known about their feeding mechanism (Jones, 1985). While their anatomical diversity has received some attention (James, 1984; Jones, 1985; Jing, 1997; Seah et al., 2009; Chakrabarty et al., 2011b), no study has quantified the degree of variation in protrusion directionality among ponyfish species or examined the morphological differences underlying this variation.

Here, we determine the anatomical basis of protrusion directionality in leiognathids. We used clearing and staining, a procedure employed for visualizing the skeletal anatomy of small vertebrates, to prepare specimens of 20 species, spanning the range of jaw protrusion direction in ponyfishes. We then photographed the cleared and stained specimens and digitally measured craniofacial morphology. This effort allowed us to describe key anatomical features associated with protrusion direction and to explore other major axes of morphological diversity within a family that exhibits extensive craniofacial variation.

2. MATERIAL AND METHODS

2.1 Study specimens & anatomical measurements

We examined 49 specimens representing 20 leiognathid species. Specimens were obtained from the Louisiana State University Museum of Natural Science and the Smithsonian National Museum of Natural History. Specimens were fixed in formalin and stored in 70% ethanol prior to this study (supplementary online material, Table 1). Each specimen was cleared and stained using procedures adapted from Dingerkus & Uhler (1977) and Taylor & Van Dyke (1985). Cleared and stained specimens were stored in glycerin and the left side of each specimen was photographed according to procedures outlined in *Photographing and Measuring Cleared and Stained Specimens* (supplementary online material). We photographed specimens in a relaxed, mostly retracted posture (Figure 1, left panel; Figure 2) and after gently manipulating the jaws into a protruded position by abducting the mandible while fixing the position of the neurocranium (Figure 1, right panel). In the absence of published video data of ponyfish feeding, we guided these manipulations based on experience with a wide diversity of percomorph fishes for which we have collected videos and manipulated specimens (Oufiero et al., 2012; Longo et al., 2016; Martinez et al., 2018).

Using NIH ImageJ software (Schneider et al., 2012), we made linear and angular measurements of morphological traits from the photographs to characterize oral jaw and craniofacial anatomy when the jaws were in the relaxed, closed mouth position (Figure 3). Linear measurements included standard length, premaxillary ascending process length (a), mandible length (b), maxilla length (c), and the vertical (d) and horizontal (e) positions of the articular-quadrate jaw joint (Figure 3A). The vertical position of the jaw joint was measured as the vertical distance between the jaw joint and the specimen's mid-horizontal axis (HA), which was defined as a line from the center of the orbit to the center of the caudal peduncle. The horizontal position of the jaw joint was measured as the horizontal distance between the jaw joint and a vertical axis (VA), drawn perpendicular to the horizontal axis and through the center of the orbit. Angular measurements included orientations of the premaxillary ascending process (f), mandible (g), and the maxilla (h;

Figure 3B). The orientations of the premaxillary ascending process and mandible were measured with respect to the vertical axis; the orientation of the maxilla was measured with respect to the horizontal axis.

Ponyfishes have previously been assigned to ventral, rostral, or dorsal protruding functional groups based on differences between the 'at rest' and 'open' mouth positions, diet, and tooth shape (Jones, 1985). We contrasted these groupings with our measure of protrusion angle, defined as premaxillary ascending process orientation relative to the vertical axis when specimens were in the opened-mouth, protruded jaw position. This measurement confirmed that our sample of species fell into three non-overlapping functional groups: ventral protruders with a protrusion angle of 28°- 39°; rostral protruders with a protrusion angle of 53°- 57°; and dorsal protruders with a protruders with a protruders, six are rostral protruders, and three are dorsal protruders resulting in ~50% sampling fraction for each functional group based on previous assignments (Jones, 1985).

2.2 Anatomical and statistical comparisons of oral jaw functional groups

We computed species means for each morphological trait and log transformed mean values for linear traits to better approximate a normal distribution of the data in R v. 3.5.1 (R Core Team, 2019). Using phylogenetic regression, we size corrected trait measurements that showed a significant relationship with standard length, including ascending process length, mandible length, maxilla length, and vertical position of the jaw joint. This correction was performed using packages *ape* (Paradis et al., 2018), *phytools* (Revell, 2018), and *geiger* (Harmon et al., 2019) and a trimmed version of the most comprehensive, time-calibrated ponyfish phylogeny

(Chakrabarty et al., 2011b). For each species in the dataset, we captured residuals with the *phyl.resid* function under a Brownian Motion model for each size dependent trait. All eight measurements were then converted to standard normal deviates (i.e., z-scores) using the *scale* function (Becker et al., 1988) to ensure that all measurements were in commensurate units and scales, thus producing a Euclidean morphospace for subsequent multivariate analyses (Huttegger & Mitteroecker, 2011; Adams & Collyer, 2019). These values were used for all subsequent analyses.

To visualize the multivariate morphological data, we performed a Principal Component Analysis (PCA) on the correlation matrix of all eight traits using the *prcomp* function. The resulting PC loadings were examined to determine how each morphological trait contributes to the variation along each PC axis. We used phylogenetic regressions to explore the pairwise relationships between traits and examined violin plots to explore trait variation within individual ponyfish functional groups. To test whether the functional groups have statistically different jaw and craniofacial morphology, we performed a phylogenetic multivariate analysis of variance (phylogenetic MANOVA) on the eight morphological traits over 10,000 iterations to test for statistical significance under a Brownian Motion model. This same method, implemented with the procD.pgls function (Adams, 2014; Collyer et al., 2015; Adams & Collyer, 2015, 2016, and 2018a) within the geomorph package (Adams et al., 2019), was used to perform individual phylogenetic ANOVAs on each of the eight traits. Finally, we performed pairwise comparisons of mean trait values among functional groups (i.e., ventral-rostral protruders, ventral-dorsal protruders, and rostral-dorsal protruders) using the *pairwise* function (Collyer et al., 2015; Adams & Collyer, 2018b) in the RRPP package (Adams & Collyer, 2020). Resulting, pairwise zvalues and p-values were evaluated to determine the magnitude and significance of the Euclidean distance between mean trait values of each functional group contrast, respectively.

3. RESULTS

3.1 Significant anatomical differentiation of functional groups

The first two axes of the PCA show some separation of functional groups in morphospace with PCs 1 and 2 describing 87.6% of variation in the data (Figure 5; Table 1). Ventral, rostral, and dorsal protruders group together along PC 1 from low to high values, respectively, suggesting that this axis primarily captures variation in species anatomy related to protrusion direction. A low value on this axis is associated with a long ascending process, a posteriorly positioned jaw joint, and the maxilla oriented with its distal arm directed posteriorly, while species with a positive position on PC 1 have a short ascending process, anteriorly and more dorsally positioned jaw joint, a more horizontally oriented ascending process, and an upright mandible (Figure 5). A phylogenetic regression of PC 1 scores against the standard normal deviates of protrusion angle confirms a strong relationship between this primary axis of morphological variation and our functional categorization of protrusion direction ($p = 1.33^{-06}$; Figure 6A). Furthermore, individual phylogenetic regressions of protrusion angle against each of our eight traits reveal significant relationships with the horizontal position of the jaw joint ($p = 1.04^{-02}$; Figure 7A), the vertical position of the jaw joint ($p = 7.05^{-03}$; not pictured), premaxillary ascending process length (p =1.54⁻⁰³; not pictured), premaxillary ascending process orientation ($p = 5.71^{-07}$; not pictured), mandible orientation ($p = 5.80^{-03}$; Figure 7B), and maxilla orientation ($p = 1.51^{-04}$; Figure 7C).

The lengths of the mandible and maxilla are strongly and positively correlated with PC 2, where $p = 4.47^{-14}$ and 1.24^{-11} , respectively (Figure 6B; Table 1). This PC axis separates the majority of species (n = 16) from a group of four rostral protruding species, including *Gazza minuta*, *G. rhombea*, *G. squamiventralis*, and *G. achlamys*. Indicated by the strong, positive loading of the lengths of the upper and lower jaw bones on PC 2 (Table 1), these four species have a larger jaw apparatus than other ponyfishes. This pattern suggests a secondary axis of morphological variation in the size of the jaws and mouth. Phylogenetic regressions of mandible and maxilla length against the horizontal and vertical position of the jaw joint show a significant, although weak, relationship between the vertical position of the jaw joint and maxilla length (p = 0.023). However, all other comparisons between the jaw joint position and jaw length yielded insignificant relationships. In combination with the insignificant relationship between protrusion angle and mandible length (p = 0.49; Figure 7D), these findings reinforce the PCA result that premaxillary protrusion direction (PC 1) and jaw size (PC 2) represent two independent axes of morphological variation in ponyfishes.

Violin plots suggest anatomical differentiation between oral jaw functional groups for all traits excluding mandible and maxilla length. Notably, the position of the jaw joint, premaxillary ascending process length, and all angles of orientation distinguish the dorsal protruders from the ventral and rostral protruders as their trait value distributions show little to no overlap (Figure 8). High variance in mandible and maxilla length of rostral protruders reflects the diversity of rostral protruding species along PC2 (Figure 5). Finally, violin plots show greater morphological variance among ventral protruding species (Figure 8), which is also apparent in the PCA as ventral protruding species occupy more area in the morphospace compared to rostral and dorsal protruding species (Figure 5).

The phylogenetic MANOVA shows a significant overall difference between functional groups (p $= 1.00^{-4}$). Phylogenetic ANOVAs of individual traits reveal significant differences between functional group means for all variables except length of the maxilla (i.e., seven of the eight traits; Table 2). Pairwise comparisons of functional groups for each trait reveal that ventral and dorsal protruding species are the most morphologically different from each other, showing significant differences for all traits excluding mandible and maxilla length. Rostral and dorsal protruders differ significantly in four traits, including premaxillary ascending process length and orientation, maxilla orientation, and vertical position of the jaw joint. Ventral and rostral protruders only differ significantly in mean trait values for mandible length and orientation, as well as premaxillary ascending process orientation. Effect size results (i.e., pairwise z-values) reveal which traits are most statistically important in differentiating between functional groups as this analysis quantifies the magnitude of difference between mean trait values (Table 2). Here, all traits that had significant p-values (p ≤ 0.05) also had large effect sizes where $z \geq |2|$, indicating that functional group mean trait values are separated by at least two standard deviations. Premaxillary ascending process length and orientation, as well as maxilla orientation most distinguish ventral and dorsal protruders. Premaxillary ascending process length, maxilla orientation, and the vertical position of the jaw joint show the largest effect sizes between rostral and dorsal protruding species. Ventral and rostral protruding species show the strongest differences in mandible length and orientation, as well as premaxillary ascending process orientation.

4. DISCUSSION

There is little known about the anatomical basis of jaw protrusion directionality despite considerable variation among extant fishes. While most suction feeding acanthomorph fishes open their mouth and protrude their jaws in a rostral direction, a number of demersal lineages that feed on benthic prey, including many sciaenids, mullids, callionymids, and gerreids, have ventrally protruding jaws. In this study, we explored the craniofacial anatomy of ponyfishes, a family exhibiting unusual variation in premaxillary protrusion orientation. Our anatomical comparison of the three oral jaw functional groups within Leiognathidae broadens our general understanding of the relationship between feeding morphology and premaxillary protrusion directionality during feeding. Overall, significant anatomical differentiation of functional groups suggested that the three angles of orientation, premaxillary ascending process length, and the position of the jaw joint discriminate the three oral jaw functional groups.

Despite extensive variation in protrusion directionality, the overall shape and orientation of the anatomical components examined herein suggest that the jaw protrusion mechanism is fairly similar across this group. Specifically, our observations and manipulations of cleared and stained specimens indicate that ponyfish protrusion involves a common perciform mechanism that is actuated by lower jaw depression (Schaeffer & Rosen, 1961; Alexander, 1967b; Liem, 1970). Ligamentous connections between the coronoid process of the mandible and the distal end of the maxillary shaft results in the distal region of the maxilla being rotated anteriorly as the mandible is depressed. As the maxilla rotates, the deep process on its proximal head presses against the ventral surface of the ascending process of the premaxilla. This pressure is resisted by 1) a

ligamentous loop that passes over the ascending process which connects the right and left maxillary heads and 2) the posteriorly tapered shape of the ascending process with associated rostral cartilage. Thus, the premaxilla responds to this maxillary pressure by sliding anteriorly into the protruded position. This basic mechanism appears to operate in all of the species we studied.

How then do the anatomical differences translate to variation in the direction of jaw protrusion in ponyfishes? One key appears to be the position of the articular-quadrate joint. The anterior position of the joint in dorsal protruders results in an upright, nearly vertical position of the adducted mandible. A more posterior position of the jaw joint in ventral protruding species results in a nearly horizontal orientation of the adducted mandible (Figure 1). In our cleared and stained specimens, a moderate mandible abduction of about twenty degrees results in a dorsally oriented mandible in the dorsal protruders, but a distinctly ventral orientation in the ventral protruders. In contrast to ventral and dorsal protruding species, rostral protruders have an intermediate jaw joint position and orientation of the resting mandible. Thus, the position of the jaw joint along the anterior-posterior axis of the fish strongly affects the resting orientation of the mandible, setting the boundary for mandibular rotation and, in turn, the protrusion angle of the upper jaw.

While the position of the jaw joint appears to play a key role in determining protrusion directionality, other variables contribute to the strong relationship between PC1 and protrusion direction (Figure 6A; Table 1). The distal end of the maxillary arm is more anteriorly oriented at rest in dorsal protruders and posteriorly oriented in ventral protruders (Figures 1, 2, and 7c).

Also, the premaxillary ascending process is more vertically oriented in the ventral protruders and almost horizontally in dorsal protruders at rest (Figures 1 and 2). Finally, ventral protruders have the longest ascending process and dorsal protruders the shortest (Figures 1, 2, and 8).

Our results also reveal that jaw and mouth size represent a second major axis of variation among ponyfish species. Our findings show that PC1 correlates strongly with variation in protrusion directionality while PC 2 primarily reflects differences in jaw size, indicating that these traits represent independent axes of diversity in this group. As proxies for jaw size, maxilla and mandible length show considerable variation within ventral and rostral protruding species but have little effect on protrusion directionality (Figure 7D). The length of the maxilla, length of the mandible, and location along the mandible where these two bones are connected form a large portion of the circumference of the gape when the jaws are extended. Thus, we propose that changes to this combination of traits results in differences in overall gape size.

Variation in protrusion directionality and jaw size in leiognathids is likely associated with differences in feeding ecology similar to the ecomorphological patterns seen in other fish groups (Liem, 1967; Chao & Musick, 1977; Liem, 1979; Gosline, 1984). Though ponyfish diets are taxonomically diverse and vary among species, some patterns emerge with respect to protrusion directionality. The diet of ventral protruding species comprises benthic prey, including infaunal polychaetes, molluscs, harpacticoid copepods, and a substantial contribution of planktonic calanoid copepods, crab larvae, and other midwater prey (Tham, 1950; Kuthalingham, 1958; Tiews et al., 1968; Hajisamae et al., 2004; Kanou et al., 2004). Some rostral protruding species have been found to feed on other fish, polychaetes, and copepods (Blaber, 1980; Jing, 1997;

Seah et al., 2009). Dorsal protruding species are recorded as primarily feeding on small midwater plankton, including calanoid copepods, and fish eggs (Venkataraman, 1960; Tiews et al., 1968; Blaber, 1980). Thus, while the diversity of feeding habits among ponyfishes deserves additional attention, a substantial literature does indicate both extensive dietary overlap among species (Tiews et al., 1968; Seah et al., 2009), as well as a tendency for ventral, rostral, and dorsal protruders to feed at progressively more elevated positions in the water column (Jing, 1997; Jones, 1985).

5. CONCLUSIONS

Unusual diversity of protrusion directionality among ponyfishes is based on variation in the location of the articular-quadrate jaw joint, combined with the size and positioning of the upper and lower jaw bones. The overall size of the jaws, indicated by length of the mandible and maxilla, varies independently of the direction of protrusion. Our study suggests two important areas for future investigation. First, our estimates of jaw function, especially directionality of protrusion, should be confirmed with kinematic analyses of live feeding videos. Second, while ponyfishes show unusually wide variation in protrusion angle among closely related species, there is considerable diversity in this trait across acanthomorph and cypriniform fishes. A key goal of future research will be to determine whether variation in protrusion direction is commonly associated with the same anatomical features found in ponyfishes. Anterior-posterior variation of the position of the articular-quadrate jaw joint may be a widespread anatomical mechanism for adjusting the directionality of jaw protrusion in teleosts, but it remains to be seen whether other groups have evolved different traits to facilitate this underappreciated aspect of diversity in fish feeding functional morphology.

Author contributions: Alexus Roberts: Conceptualization; methodology; investigation; data curation; formal analysis; visualization; writing – original draft; writing – review & editing; funding acquisition. Jennifer Hodge: Methodology; investigation; formal analysis – supporting; writing – review & editing. Prosanta Chakrabarty: Resources; writing – review & editing. Peter Wainwright: Conceptualization; resources; formal analysis – supporting; writing – original draft; writing – review & editing; funding acquisition.

Acknowledgments: We thank K. Corn, S. Friedman, C. Martinez, E. Burress, A. Barley, and D. Satterfield for critical input; M. Rupp for assistance with clearing and staining specimens; and D. Adams and M. Collyer for advice on data treatment. S. Parker of the Louisiana State University Museum of Natural Science and J. Williams of the Smithsonian National Museum of Natural History generously facilitated specimen loans. This research was supported by the Center for Population Biology at UC Davis and the National Science Foundation supported A.S.R. with a Graduate Research Fellowship under Grant No. 1650042, J.R.H. with a Postdoctoral Research Fellowship in Biology for Research using Biological Collections (DBI-1523934), and grant DEB-1061981 to P.C.W.

Data Availability: The data that supports the findings of this study are available in the supplementary material of this article.

Conflict of Interest: The authors have no conflicts of interest to report.

References:

Adams, A. D., & Collyer, M. L. (2020). Package ' RRPP '.

Adams, A. D., Collyer, M. L., & Kaliontzopoulou, A. (2019). Package ' geomorph'.

- Adams, D. C. (2014). A method for assessing phylogenetic least squares models for shape and other high-dimensional multivariate data. *Evolution*, 68, 2675-2688. <u>https://doi.org/10.1111/evo.12463</u>
- Adams, D. C., & Collyer, M. L. (2015). Permutation tests for phylogenetic comparative analyses of high-dimensional shape data: what you shuffle matters. *Evolution*, 69, 823-829. <u>https://doi.org/10.1111/evo.12596</u>
- Adams, D. C., & Collyer, M. L. (2016). On the comparison of the strength of morphological integration across morphometric datasets. *Evolution*, 70, 2623-2631. <u>https://doi.org/10.1111/evo.13045</u>
- Adams, D. C. & Collyer, M. L. (2018a). Multivariate phylogenetic comparative methods: evaluations, comparisons, and recommendations. *Systematic Biology*, 67(1),14-31. <u>https://doi.org/10.1093/sysbio/syx055</u>
- Adams, D. C & Collyer, M. L. (2018b). Phylogenetic ANOVA: group-clade aggregation, biological challenges, and a refined permutation procedure. *Evolution*, 72(6), 1204-1215. <u>https://doi.org/10.1111/evo.13492</u>
- Adams, D. C. & Collyer, M. L. (2019). Phylogenetic methods and the evolution of multivariate phenotypes. *Annual Review of Ecology, Evolution, and Systematics* 50, 405-425. https://doi.org/10.1146/annurev-ecolsys-110218-024555
- Alexander, R. McN. (1967a). Mechanics of the jaws of some atheriniform fish. *Journal of Zoology, London*, 151, 233-255. <u>https://doi.org/10.1111/j.1469-7998.1967.tb02876.x</u>
- Alexander, R. McN. (1967b). The functions and mechanisms of the protrusible upper jaws of some acanthopterygian fish. *Journal of Zoology, London*, 151, 43-64. <u>https://doi.org/10.1111/j.1469-7998.1967.tb02865.x</u>
- Becker, R. A., Chambers, J. M., & Wilks, A. R. (1988). The New S Language: A Programming Environment for Data Analysis and Graphics. Pacific Grove, California: Wadsworth & Brooks/Cole.
- Blaber, S. J. M. (1980). Fish of the Trinity Inlet system of north Queensland with notes on the ecology of fish faunas of tropical Indo-Pacific estuaries. *Australian Journal of Marine* and Freshwater Research, 31(2), 137-146. <u>https://doi.org/10.1071/MF9800137</u>
- Chakrabarty, P., Davis, M.P., Berquist, R., Gledhill, K., Sparks, J., & Frank, L. (2011a) Evolution of the light organ system in ponyfishes (Teleostei: Leiognathidae). *Journal of Morphology*, 272: 704-721. <u>https://doi.org/10.1002/jmor.10941</u>

- Chakrabarty, P., Davis, M. P., Smith, W. L., Baldwin, Z. H., & Sparks, J. S. (2011b). Is sexual selection driving diversification of the bioluminescent ponyfishes (Teleostei: Leiognathidae)? *Molecular Ecology*, 20(13), 2818-2834. <u>https://doi.org/10.1111/j.1365-294X.2011.05112.x</u>
- Chao, L. N., & Musick, J. A. (1977). Life-history, feeding-habits, and functional-morphology of juvenile sciaenid fishes in the York River estuary, Virginia. *Fishery Bulletin*, 7(4), 657-702. <u>https://scholarworks.wm.edu/vimsarticles/637</u>
- Collyer, M. L., Sekora, D. J., & Adams, D. C. (2015). A method for analysis of phenotypic change for phenotypes described by high-dimensional data. *Heredity*, 115, 357-365. <u>https://doi.org/10.1038/hdy.2014.75</u>
- Dingerkus, G., & Uhler, L. D. (1977). Enzyme clearing of alcian blue stained whole small vertebrates for demonstration of cartilage. *Stain Technology*, 52(4), 229-232. https://doi.org/10.3109/10520297709116780
- Gibb, A. C., Staab, K., Moran, C., Ferry, L.A. (2015). The teleost intramandibular joint: a mechanism that allows fish to obtain prey unavailable to suction feeders. *Integrative and Comparative Biology*, 55(1), 85-96. <u>https://doi.org/10.1093/icb/icv042</u>
- Gosline, W. A. (1980). The evolution of some structural systems with reference to the interrelationships of modern lower teleostean fish groups. *Japanese Journal of Ichthyology*, 27(1), 1-27. <u>https://doi.org/10.11369/jji1950.27.1</u>
- Gosline, W. A. (1984). Structure, function, and ecology in the Goatfishes (family Mullidae). *Pacific Science*, 38(4), 312-323. <u>http://hdl.handle.net/10125/832</u>
- Grobecker, D. B. (1983). The 'lie-in-wait' feeding mode of a cryptic teleost, *Synanceia verrucose*. In, D. L. G. Noakes, D. G., Lindquist, G. S., Helfman, & J. A., Ward (Eds.), *Predators and Prey in Fishes* (pp. 29-40). Dordercht, The Netherlands: Springer.
- Hajisamae, S., Chou, L. M., & Ibrahim, S. (2004). Feeding habits and trophic relationships of fishes utilizing an impacted coastal habitat, Singapore. *Hydrobiologia*, 520, 61-71. <u>https://doi.org/10.1023/B:HYDR.0000027727.90956.a9</u>

Harmon, A. L. et al. (2019). Package ' geiger '.

- Harvey, E. N. (1921). A fish, with a luminous organ, designed for the growth of luminous bacteria. *Science*, 53, 314-315. https://doi.org/10.1126/science.53.1370.314
- Hastings, J. W. (1971). Light to hide by: Ventral luminescence to camouflage the silhouette. *Science*, 173, 1016-1017. <u>https://doi.org/10.1126/science.173.4001.1016</u>
- Holzman, R., Day, S. W., & Wainwright, P. C. (2007). Timing is everything: coordination of strike kinematics affects the force exerted by suction feeding fish on attached prey. *Journal of Experimental Biology*, 210, 3328-3336. <u>https://doi.org/10.1242/jeb.008292</u>

- Accepted Article
- Huttegger, S. M. & Mitteroecker, P. (2011). Invariance and meaningfulness in phenotypic spaces. *Evolutionary Biology*, 38, 335-351. <u>https://doi.org/10.1007/s11692-011-9123-x</u>
- James, P. S. B. R. (1984). Leiognathidae. In, W. Fischer and G. Bianchi (Eds.), FAO species identification sheets for fishery purposes, Western Indian Ocean, Fishing Area 51 (Vol. 2.). FAO, Rome.
- Jing, L. (1997). Morphological comparison of the feeding apparatus of three co-occurring species of ponyfishes (Leiognathidae). *Chinese Journal of Oceanology and Limnology*, 15(3), 224-226. <u>https://link.springer.com/content/pdf/10.1007/BF02850877.pdf</u>
- Jones, G. (1985). Revision of the Australian species of the fish family Leiognathidae. *Australian Journal of Marine and Freshwater Research*, 36(4), 559-613. <u>https://doi.org/10.1071/MF9850559</u>
- Kanou, K., Sano, M., & Kohno, H. (2004). Food habits of fishes on unvegetated tidal mudflats in Tokyo Bay, central Japan. *Fisheries Science*, 70(6), 978-987. <u>https://doi.org/10.1111/j.0919-9268.2004.00897.x</u>
- Konow, N. & Bellwood, D. R. (2005). Prey-capture in *Pomacanthus semicirculatus* (Teleostei, Pomacanthidae): functional implications of intramandibular joints in marine angelfishes. *Journal of Experimental Biology*, 208, 1421-1433. <u>https://doi.org/10.1242/jeb.01552</u>
- Kuthalingham, M. (1958). The food and feeding habits of some young Silver-bellies. *Journal of the Madras University*, 28, 13-22.
- Lauder, G. V. (1980). Hydrodynamics of prey capture by teleost fishes. In, Schneck, D. J. (Eds.), *Biofluid Mechanics 2* (pp. 161-181). New York: Plenum Press. <u>https://doi.org/10.1007/978-1-4757-4610-5_9</u>
- Liem, K. F. (1967). Functional morphology of the head of the anabantoid teleost fish *Helostoma temmincki*. Journal of Morphology, 121,135-158. <u>https://doi.org/10.1002/jmor.1051210204</u>
- Liem, K. F. (1970). Comparative functional anatomy of the Nandidae (Pisces: Teleostei). Chicago: Field Museum of Natural History. <u>https://doi.org/10.5962/bhl.title.3081</u>
- Liem, K. F. (1979). Modulatory multiplicity in the feeding mechanism in cichlid fishes, as exemplified by the invertebrate pickers of Lake Tanganyika. *Journal of Zoology, London*, 189, 93-125. <u>https://doi.org/10.1111/j.1469-7998.1979.tb03954.x</u>
- Liem, K. F. (1980). Acquisition of energy by teleosts: Adaptive mechanisms and evolutionary patterns. In, Ali, M. A. (Eds.), *Environmental Physiology of Fishes*, NATO Advanced Study Institute Series (Series A: Life Science), vol 35 (pp. 299-334). Boston, Massachusetts: Springer. <u>https://doi.org/10.1007/978-1-4899-3659-2_10</u>

- Longo, S. J., McGee, M. D., Oufiero, C. E., Waltzek, T. B., & Wainwright, P. C. (2016). Ram, not suction, is the primary axis of suction feeding diversity. *Journal of Experimental Biology*, 219, 119-128. <u>https://doi.org/10.1242/jeb.129015</u>
- Martinez, C. M., McGee, M. D., Borstein, S. R., & Wainwright, P. C. (2018). Feeding ecology underlies the evolution of cichlid jaw mobility. *Evolution*, 72(8), 1645-1655. <u>https://doi.org/10.1111/evo.13518</u>
- McFall-Ngai, M. J., & Dunlap, P. V. (1984). External and internal sexual dimorphism in leiognathid fishes: Morphological evidence for sex-specific bioluminescent signaling. *Journal of Morphology*, 182(1), 71-83. <u>https://doi.org/10.1002/jmor.1051820105</u>
- Motta P. J. (1984). Mechanics and functions of jaw protrusion in teleost fishes—a review. *Copeia*, 1984(1) 1–18. <u>https://doi.org/10.2307/1445030</u>
- Muller, M. & Osse, J. W. M. (1978). Structural adaptation to suction feeding in fish. In, *Proceedings of the Zodiac Symposium on Adaptation* (pp. 57-60). Wageningen, Netherlands: Pudoc.
- Nelson, J. S., Grande, T. C., & Wilson, M. V. H. (2016). Fishes of the World (Fifth Edition). Hoboken: John Wiley & Sons. <u>https://doi.org/10.1002/9781119174844</u>
- Oufiero, C. E., Holzman, R. A., Young, F. A., & Wainwright, P. C. (2012). New insights from serranid fishes on the role of trade-offs in suction feeding diversification. *Journal of Experimental Biology*, 215, 3845-3855. <u>https://doi.org/10.1242/jeb.074849</u>

Paradis, E., et al. (2018). Package 'ape'.

- Pietsch, T. W. (1978). The feeding mechanism of *Stylephorus chordatus* (Teleostei: Lampridiformes): functional and ecological implications. *Copeia*, 2, 255-262. https://doi.org/10.2307/1443560
- R Core Team (2019). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/

Revell, M. L. J. (2018). Package ' phytools '.

- Sasaki, A., Ikejima, K., Aoki, S., Azuma, N., Kashimura, N., & Wada, M. (2003). Field evidence for bioluminescent signaling in the pony fish, *Leiognathus elongatus*. *Environmental Biology of Fishes*, 66, 307–311. <u>https://doi.org/10.1023/A:1023959123422</u>
- Schaeffer, B. & Rosen, D. E. (1961). Major adaptive levels in the evolution of the actinopterygian feeding mechanism. *American Zoologist*, 1, 187-204. <u>http://www.jstor.org/stable/3881250</u>
- Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image analysis. *Nature Methods*, 9, 671–675. <u>https://doi.org/10.1038/nmeth.2089</u>

- Seah, Y. G., Abdullah, S., Zaidi, C. C., & Mazlan, A.G. (2009). Systematic accounts and some aspects of feeding and reproductive biology of ponyfishes (Perciformes: Leiognathidae). *Sains Malaysiana*, 38(1), 47-56.
- Taylor, W. R. & Van Dyke, G. C. (1985). Revised procedures for staining and clearing small Fishes and other vertebrates for bone and cartilage study. *Cybium*, 9, 107-119.
- Tham, A. K. (1950). The food and feeding relationship of the fishes of Singapore Straits. In, Colonial Office Fishery Publication (pp. 1-35). H.M. Stationery Office.
- Tiews, K., Divino, P., Ronquillo, I. A., & Marquez, J. (1968). On the food and feeding habits of eight species of Leiognathus found in Manila Bay and San Miguel Bay. *Indo-Pacific Fisheries Council*, 13(3), 93-99.
- Venkataraman, G. (1960). Studies on the food and feeding relationships of the inshore fishes off Calicut on the Malabar coast. *Indian Journal of Fisheries*, 7(2), 275-306. <u>http://eprints.cmfri.org.in/id/eprint/1898</u>
- Wainwright, P. C., McGee, M. D., Longo, S. J. & Hernandez, L. P. (2015). Origins, innovations, and diversification of suction feeding in vertebrates. *Integrative and Comparative Biology*, 55(1), 134-145. <u>https://doi.org/10.1093/icb/icv026</u>
- Woodland, D. J., Cabanban, A. S., Taylor, V. M., & Taylor, R. J. (2002). A synchronized rhythmic flashing light display by schooling *Leiognathus splendens* (Leiognathidae : Perciformes). *Marine and Freshwater Research*, 53, 159–162. <u>https://doi.org/10.1071/MF01157</u>

Tables

Table 1. Principal component loadings on the first 5 PC axes. PC loadings indicate the contribution of each variable to the principal component. Bolded values represent loadings that are greater than or equal to |0.3|.

Morphological trait	Principal component loadings						
	PC1	PC2	PC3	PC4	PC5		
Ascending Process Length	-0.40	0.29	-0.02	0.41	0.14		
Mandible Length	0.03	0.60	0.09	0.16	-0.25		
Maxilla Length	0.02	0.59	0.24	-0.16	-0.20		
Ascending Process Orientation	0.43	0.22	-0.22	0.02	-0.34		
Mandible Orientation	-0.33	-0.28	0.71	0.07	-0.48		
Maxilla Orientation	0.44	0.03	0.43	-0.52	0.18		
Horizontal Jaw Joint	0.43	0.02	0.42	0.62	0.45		
Vertical Jaw Joint	-0.40	0.29	0.13	-0.36	0.54		
Percent Variance	54.1%	33.5%	6.9%	2.4%	1.6%		
Cumulative Variance	54.1%	87.6%	94.5%	96.9%	98.5%		

Table 2. Phylogenetic analyses of variance. Phylogenetic ANOVAs were performed on the standard normal deviates of the eight morphological traits. Pairwise effect sizes (i.e., z-values) indicate the magnitude of the difference between functional group means, where values $\geq |2|$ indicate that group means are separated by two or more standard deviations. Pairwise p-values indicate functional groups that exhibit significantly different morphologies for a specified trait. Significant p-values (≤ 0.05) are bolded.

T. 14	Phylogenetic ANOVA						
l rait	F-value (df = 2,18)	p-value	Functional group comparison	Pairwise z-value	Pairwise p-value		
Ascending Process Length	-		Ventral – Rostral	-1.35	0.983		
	9.27	0.002	Ventral – Dorsal	3.66	0.003		
			Rostral – Dorsal	3.72	0.002		
Mandible Length	4.29	0.029	Ventral – Rostral	2.82	0.010		
			Ventral – Dorsal	-0.68	0.694		
			Rostral – Dorsal	1.10	0.146		
Maxilla Length	2.11	0.146	Ventral – Rostral	1.89	0.054		
			Ventral – Dorsal	-0.81	0.746		
			Rostral – Dorsal	0.49	0.290		
Ascending Process Orientation	29.95		Ventral – Rostral	2.23	0.018		
		1.00 e-4	Ventral – Dorsal	4.96	1.00 e-4		
			Rostral – Dorsal	2.31	0.027		
Mandible Orientation	8.32	0.003	Ventral – Rostral	2.25	0.028		
			Ventral – Dorsal	3.48	0.002		
			Rostral – Dorsal	0.91	0.179		
Maxilla Orientation			Ventral – Rostral	-0.02	0.440		
	9.76	0.003	Ventral – Dorsal	4.06	9.00 e-4		
			Rostral – Dorsal	3.12	0.006		
Horizontal Position of the jaw joint			Ventral – Rostral	1.04	0.163		
	3.91	0.036	Ventral – Dorsal	2.74	0.009		
			Rostral – Dorsal	0.98	0.176		
Vertical Position of the jaw joint	4.80	0.020	Ventral – Rostral	-0.34	0.560		
			Ventral – Dorsal	3.04	0.006		
			Rostral – Dorsal	2.34	0.021		

Figure Legends

Figure 1: Leiognathids show diversity in the orientation of oral jaw protrusion, with species that protrude their jaws in a ventral (**A**), rostral (**B**), or dorsal (**C**) direction. Left and right panels depicts specimens in a relaxed, mostly retracted posture and a protruded jaw position, respectivley. Black bars indicate a 10 mm scale. Illustrations by A. Roberts.

Figure 2. Cleared and stained ponyfishes depict variation in craniofacial morphology and body form. Species include *Gazza achlamys* (**A**), *Secutor insidiator* (**B**), *Photopectoralis aureus* (**C**), *Leiognathus robustus* (**D**), *Eubleekeria jonesi* (**E**), and *Nuchequula decorus* (**F**). Black bars indicate a 10 mm scale.

Figure 3. Measurements made on ponyfish jaw morphology, depicted with *Leiognathus robustus*. **A**) Ascending process length (a), mandible length (b), maxilla length (c), and the vertical (d) and horizontal (e) positions of the jaw joint were measured digitally from photographs. **B**) Angular orientations of the ascending process (f; in reference to the vertical axis), mandible (g; in reference to the vertical axis), and the maxilla (h; in reference to the horizontal axis) were measured from specimen photographs in the 'at rest' mouth position. Dotted gray lines represent the horizontal (HA) and vertical (VA) axes.

Figure 4. Jaw protrusion orientation in ponyfishes. **A)** Phenogram of protrusion angle shows the clustering of dorsal, rostral and ventral protruding species. **B)** Annotated phylogeny depicts the relationships between the 20 leiognathid species examined in this study. Numbers indicate species shown in right-hand images. White bars indicate a 10mm scale.

Figure 5. A phylomorphospace projection of ponyfish species based on eight measurements of jaw and cranial anatomy. Here, each point represents a species' standard normal deviation value, with ventral, rostral, and dorsal protruding species in gray, blue, and orange, respectively. Specimen images of outlier species are outlined according their functional group. PC loadings are reported in Table 1.

Figure 6. Oral jaw protrusion direction and jaw size represent two major axes of morphological diversity in ponyfishes. Phylogenetic regression analyses of standard normal deviates reveal significant relationships between (**A**) protrusion angle and PC 1 ($p = 1.33^{-06}$; adjusted $R^2 = 0.72$) and (**B**) mandible length and PC 2 ($p = 4.47^{-14}$; adjusted $R^2 = 0.96$).

Figure 7. Phylogenetic regressions reveal significant relationships between protrusion angle and individual measurements of the jaw bones. Here, each point represents a species' standard normal deviation value with ventral, rostral, and dorsal protruding species in gray, blue, and orange, respectively. (**A**) horizontal position of the jaw joint ($p = 1.04^{-02}$; adjusted $R^2 = 0.27$), (**B**) mandible orientation ($p = 5.80^{-03}$; adjusted $R^2 = 0.32$), and (**C**) maxilla orientation ($p = 1.51^{-04}$; adjusted $R^2 = 0.53$) show a significant relationship with protrusion angle. In contrast, (**D**) protrusion angle and mandible length show a non-significant relationship (p = 0.49; adjusted $R^2 = -0.03$), with ventral and rostral protruding species exhibiting considerable variation in mandible length.

Figure 8. Violin plots show non-overlapping distributions for several morphological traits. For each trait, three distributions are shown representing the morphological variance exhibited by each functional group. Each distribution has three horizontal lines indicating the mean, upper, and lower quartiles. Each trait plot uses standard normal deviates collected from specimens in the 'at rest' mouth position.







This article is protected by copyright. All rights reserved.

10mm











Horizonatal Jaw Joint Position



Vertical Jaw Joint Position



Ascending Process Orientation



Mandible Orientation



Maxilla Orientation



Functional Group